

MEMORANDUM

November 1, 2016

Re: ICD-10 for presumptive UDT testing

Dear Provider,

Thank you for your confidence in Advanta Analytical Laboratories. The following ICD-10 codes are approved by CMS as covered for immunoassay (presumptive) screening (CPT G0479). Any carrier that utilizes HCPCS for toxicology coding/billing will follow this local coverage determination (LCD) guideline unless they have adopted an internal policy that states otherwise.

NOTE:

1. The contractor will consider **PRESUMPTIVE UDT** testing in excess of 12 per Calendar year not reasonable and necessary.
2. An exception to the above limitation will be made when patients have documented diagnoses consistent with a substance abuse disorder (SUD), for which patients **PRESUMPTIVE UDT** shall not occur more than 3 times within a seven-day period, based upon the following guidelines for monitoring abstinence.
3. For patients with 0 to 30 consecutive days of abstinence, **PRESUMPTIVE UDT** is expected at a frequency of 1 to 3 **PRESUMPTIVE UDT** per week. More than 3 **PRESUMPTIVE** panels in one week is not reasonable and necessary and is not covered by Medicare.
4. For patients with 31 to 90 consecutive days of abstinence, **PRESUMPTIVE UDT** is expected at a frequency of 1 to 3 **UDT** per week. More than 3 **PRESUMPTIVE UDT** in one week is not reasonable and necessary and is not be covered by Medicare.
5. For patients with > 90 consecutive days of abstinence, **PRESUMPTIVE UDT** is expected at a frequency of 1 to 3 **UDT** in one month. More than 3 physician-directed **UDT** in one month is not reasonable and necessary and is not covered

Primary ICD-10 codes for presumptive UDT

E87.2	Acidosis
F10.120*	Alcohol abuse with intoxication, uncomplicated
F11.20*	Opioid dependence, uncomplicated
F12.120*	Cannabis abuse with intoxication, uncomplicated
F12.220*	Cannabis dependence with intoxication, uncomplicated
F13.120*	Sedative, hypnotic or anxiolytic abuse with intoxication, uncomplicated
F14.120*	Cocaine abuse with intoxication, uncomplicated
F14.220*	Cocaine dependence with intoxication, uncomplicated
F16.120*	Hallucinogen abuse with intoxication, uncomplicated
F18.10*	Inhalant abuse, uncomplicated
F18.120*	Inhalant abuse with intoxication, uncomplicated
F18.90*	Inhalant use, unspecified, uncomplicated
F19.20*	Other psychoactive substance dependence, uncomplicated
F20.0	Paranoid schizophrenia
F20.1	Disorganized schizophrenia
F20.2	Catatonic schizophrenia
F20.89	Other schizophrenia
F55.3*	Abuse of steroids or hormones
F55.4*	Abuse of vitamins
F55.8*	Abuse of other non-psychoactive substances
G40.301	Generalized idiopathic epilepsy and epileptic syndromes, not intractable, with status epilepticus
G40.309	Generalized idiopathic epilepsy and epileptic syndromes, not intractable, without status epilepticus

Primary ICD-10 codes for presumptive UDT (Page 2)

G40.311	Generalized idiopathic epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.319	Generalized idiopathic epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.901	Epilepsy, unspecified, not intractable, with status epilepticus
G40.909	Epilepsy, unspecified, not intractable, without status epilepticus
G40.911	Epilepsy, unspecified, intractable, with status epilepticus
G40.919	Epilepsy, unspecified, intractable, without status epilepticus
I44.0	Atrioventricular block, first degree
I44.1	Atrioventricular block, second degree
I44.30	Unspecified atrioventricular block
I45.81	Long QT syndrome
I47.1	Supraventricular tachycardia
I47.2	Ventricular tachycardia
M25.50	Pain in unspecified joint
M47.811	Spondylosis without myelopathy or radiculopathy, occipito-atlanto-axial region
M47.812	Spondylosis without myelopathy or radiculopathy, cervical region
M47.813	Spondylosis without myelopathy or radiculopathy, cervicothoracic region
M47.816	Spondylosis without myelopathy or radiculopathy, lumbar region
M47.817	Spondylosis without myelopathy or radiculopathy, lumbosacral region
M47.818	Spondylosis without myelopathy or radiculopathy, sacral and sacrococcygeal region
M51.14	Intervertebral disc disorders with radiculopathy, thoracic region
M51.15	Intervertebral disc disorders with radiculopathy, thoracolumbar region
M51.16	Intervertebral disc disorders with radiculopathy, lumbar region
M51.17	Intervertebral disc disorders with radiculopathy, lumbosacral region
M51.36	Other intervertebral disc degeneration, lumbar region

Primary ICD-10 codes for presumptive UDT (Page 3)

M51.37 Other intervertebral disc degeneration, lumbosacral region
M54.10 Radiculopathy, site unspecified
M54.14 Radiculopathy, thoracic region
M54.15 Radiculopathy, thoracolumbar region
M54.16 Radiculopathy, lumbar region
M54.17 Radiculopathy, lumbosacral region
M54.18 Radiculopathy, sacral and sacrococcygeal region
M54.2 Cervicalgia
M54.5 Low back pain
M60.88 Other myositis, other site
M60.89 Other myositis, multiple sites
M79.1 Myalgia
M79.2 Neuralgia and neuritis, unspecified
M79.7 Fibromyalgia
R40.0 Somnolence
R40.1 Stupor

Please contact Advanta with any questions or concerns.

Sincerely,



JULIE JUMALON

ADVANTA ANALYTICAL LABORATORIES

10935 CR 159, Tyler, TX 75703

OFFICE: 903.707.2197

FAX: 903.839.2494 •

jjumalon@aalabs.com